

Improving Physical Function and Performance With Progressive Resistance Strength Training in Older Adults

1. Kathleen Kline Mangione,
2. Amy H. Miller and
3. Irene V. Naughton

± Author Affiliations

1. *K.K. Mangione, PT, PhD, GCS, is Professor, Department of Physical Therapy, Arcadia University, Glenside, Pennsylvania.*
2. *A.H. Miller, PT, DPT, is Assistant Professor of Physical Therapy and Curricular Coordinator of the Entry-Level DPT Program, Department of Physical Therapy, Arcadia University, Glenside, Pennsylvania.*
3. *I.V. Naughton, PT, MS, is Team Clinical Coordinator of Physical Therapy, Abington Memorial Hospital, Home Health Care Services, Abington, Pennsylvania.*

Can progressive resistance training help her regain function so that she can function independently in the community again?

Age-related muscle loss, also known as **sarcopenia**, is a common problem among **older adults**. Physical therapists are concerned about muscle loss because it is associated with loss of muscle strength and muscle quality.² Muscle strength—the amount of force a muscle can produce—is highly associated with functional activities.³⁻⁴ Loss of strength is associated with slow gait speed, poor endurance, inability to rise from a chair, falls risk, incident disability, and frailty.⁵⁻⁷ Men and women demonstrate differing trajectories in muscle loss, with men's loss being more gradual and women showing a more dramatic decrease after menopause.³ The muscle loss of sarcopenia is a problem for both older men and women and increases with increasing age. Prevalence estimates from the National Health and Nutrition Examination Survey (NHANES) data suggest that 35% of the older adults had moderate sarcopenia, whereas 10% had severe sarcopenia.⁸ Among a sample of community-dwelling volunteers, sarcopenia occurred in 22.6% of women and 26.8% of men. In the subgroup of women and men who were 80 years and older, prevalence rates of sarcopenia were 31.0% and 52.9%, respectively.⁹

Resistance training is used to counteract the effects of muscle loss. It is a form of exercise in which the muscle contracts against an external load.¹⁰ Some of the equipment commonly used to perform resistance training includes free weights, exercise machines, body weight, and elastic bands. Progressive resistance training (PRT), in which the load is systematically increased as the person is able to work against a heavier load, has been shown to produce large increases in strength and moderate increases in bone mineral density, lean body mass, insulin sensitivity, and submaximal and maximal endurance.¹⁰ Although an initial Cochrane review showed that PRT produced large effects in improving muscle strength,¹¹ many new trials have been added to the literature. The authors note that uncertainty remains about the effectiveness of PRT on physical disability outcomes, on more pragmatic home or hospital-based programs, on older adults who have multiple health problems, and on the comparative benefits of PRT versus other exercise programs.¹² The purpose of the review by Liu and Latham¹² was to determine the effects of PRT on physical function in older adults. The primary outcome was a measure of activity and participation (activities of daily living and the physical function component of health-related quality of life

scores). The secondary outcomes were measures of body structure and function (muscle strength, aerobic capacity) and measures of activity (balance, gait speed, Timed “Up & Go” [TUG] test times, and chair rise speed). Additional outcomes were adverse events, hospital admissions, and death.

[Next Section](#)

Take-Home Message

The Cochrane review by Liu and Latham included 121 trials with more than 6,700 older adults. The results are presented as standardized mean difference (SMD), a unitless measure that represents effect size. It has been reported that effect sizes <0.2 are small, effects >0.2 to <0.8 are moderate, and effects >0.8 are large. When comparing PRT to a control group, Lin and Lantham found a small but significant effect for PRT for improving physical disability (SMD=0.14; 95% confidence interval [CI]=0.05, 0.22). A moderate to large benefit was found for PRT for improving strength (SMD=0.84; 95% CI=0.67, 1.00). Progressive resistance training improved aerobic capacity (SMD=0.31; 95% CI=0.09, 0.53). Progressive resistance training had a small positive effect on balance but the value was not significant (SMD=0.12; 95% CI=0.00, 0.25). Chair rise speed showed a large effect (speeds became faster) after PRT (SMD=-0.94; 95% CI=-1.49, -0.38). Six-minute walk distance, gait speed, TUG test values after PRT are reported in the actual units in the [Table](#).

View this table:

[In this window](#)

[In a new window](#)

[Download as PowerPoint Slide](#)

Table.

Progressive Resistance Training (PRT) for Improving Physical Function in Older Adults: Cochrane Review Results

Because of the number of studies, the variability in characteristics of the participants, the design of the PRT program, and the outcomes assessed, the authors conducted multiple subanalyses of the data. The [Table](#) highlights the subanalyses that might be most relevant to physical therapist practice. As the authors indicate, caution has to be applied when interpreting the subanalyses because the results are based on fewer studies.

The take-home message regarding progressive resistance training is clear: strength training performed 2 to 3 times per week reduces physical disability and improves some functional abilities, such as gait speed and chair rise time. Progressive resistance training has its largest effect in improving strength. There was no subgroup or analysis that showed that strength training was detrimental to older adults. Adverse events typically were not reported, but when they *were* reported, joint and muscle pain were the most common adverse events. The authors conclude that PRT is an appropriate intervention for older adults who want to improve performance of simple physical tasks.

[Previous Section](#)[Next Section](#)

Case #4: Applying Evidence to a Patient With Muscle Weakness

Can progressive resistive training help this patient?

Mrs. Smith is an 86-year-old woman who was living alone in a farmhouse when she fell on the floor and remained there for 2 days. Her admitting diagnoses to the hospital were dehydration, hypertension, and ambulatory dysfunction because of the fall. Her history also included left shoulder rotator cuff impairment, cardiomegaly, bilateral osteoarthritis, spinal stenosis, and anemia. She was referred for home health physical therapy and was discharged to her son's home with the goal of living independently in a nearby apartment. To achieve this goal, improvements needed to be made in her physical and functional abilities. Upon initial examination, Mrs. Smith appeared cognitively intact (alert and oriented to time, place, and person). Her vital signs were as follows: heart rate, 88 beats per minute; blood pressure, 122/82 mm Hg; respiration rate, 20. Her lungs were clear. Her impairments and functional limitations included decreased lower-extremity strength, impaired balance, slow gait speed, assistance needed in activities of daily living (ADLs), and fall risk. The TUG test took 53 seconds (>30 seconds suggests dependence in ADLs),¹⁶ and she required close supervision to stand from the chair. She also required supervision when climbing steps and required contact guarding when stepping over a tub. Her gait speed was 0.28 m/s (normal speed is 1.0 m/s¹⁷), and she needed contact guarding with a wheeled walker. Her Outcome and Assessment Information Set (OASIS—mandated tool by the Centers for Medicare & Medicaid Services) Falls Risk Assessment score was 14 (scores >10 suggest high risk for falls¹⁸).

How did the results of the Cochrane systematic review apply to Mrs. Smith?

Mrs. Smith was similar to the community-dwelling older adults addressed in the systematic review (Table). Even though most studies included in the review took place in a gym or clinic setting, the physical therapist and Mrs. Smith decided she would be a candidate for home-based progressive resistance training. Consistent with the frequency of training reported in the review, visits were planned for 2 times per week. The physical therapist had an adjustable cuff weight, and, because the patient was ambulatory, the therapist decided to try to use 10 lb to assess Mrs. Smith's strength. Mrs. Smith could lift 10-lb cuff weights on each leg for an 8-repetition maximum before fatigue. Thus, her 8-repetition maximum was 10 lb. The PRT dose began with 10 lb and she increased the load on subsequent visits so that Mrs. Smith would always fatigue between 6 and 10 repetitions. She performed 2 sets of the knee extension exercises with repetitions to fatigue (varied from 6 to 10 repetitions) for each set. At the second session, Mrs. Smith complained of increased quadriceps soreness and was unsure whether she should continue with PRT or not. The physical therapist explained to Mrs. Smith that muscle soreness was to be expected but would dissipate with continued PRT. Mrs. Smith agreed to continue with training.

Other interventions included in the home physical therapy visits were gait training indoors and outdoors, progressing from a wheeled walker to a standard cane indoors, stair climbing indoors and outdoors, transfer training, and standing weight shifting to improve her balance. Her home exercise program consisted of daily active range of motion exercises for the upper and lower extremities in both sitting and standing positions, along with tandem and sideways ambulation with single upper-extremity support.

How well do the outcomes of intervention provided to Mrs. Smith match those suggested by the systematic review?

Mrs. Smith received 7 home visits over the 3.5 week episode of care. The episode was shorter than anticipated because she moved out of the area. Her outcomes were consistent or greater than changes reported in the systematic review. She could lift a 20-lb weight 8 times before fatigue, indicating a strength gain of 100%. Mrs. Smith's measured strength gains exceeded those reported in the review regardless of health status or functional grouping. Her TUG score decreased by 27 seconds, now requiring 26 seconds to complete the test. This change is well over reported minimal detectable changes reported for the TUG (Table), and far more substantial than the change reported in the Cochrane review (0.69 seconds). The change in TUG time may be related more to the functional training activities that were part of Mrs. Smith's program than solely to the PRT. Mrs. Smith's gait speed increased to 0.38 m/s, a 0.1 m/s improvement that is consistent with the systematic review, which reported a 0.08 m/s improvement in gait speed following progressive resistance training. Mrs. Smith had progressed to independent indoor ambulation with a standard cane. Her transfers were now independent, which showed as a gain in the OASIS transfer measure (M1850). Mrs. Smith achieved her goal of returning to independent living in a smaller space without relying on her children for performance of ADLs. Progressive resistance training contributed to her independence in ADLs, which was the primary outcome of the Cochrane review.

Can you apply the results of the systematic review to your own patients?

The findings of this Cochrane review apply well to Mrs. Smith. The age of the participants in the studies ranged from 60 to more than 80 years. The review included studies with both healthy participants and older adults with specific health problems, as well as participants with and without functional limitations. Trials also examined older adults residing in the community or within institutions. Mrs. Smith was well represented in the review.

This Cochrane Review has some limitations, one of which is that it excluded studies that included balance, aerobic, or other training as part of the exercise intervention. In addition, only quadriceps force measures were included in the calculation of effect sizes. Only 2 studies were conducted in hospitals, which may limit application to patients in the acute hospital setting.

The common features of training noted in the systematic review were that the programs included a supervised progression of the resisted exercise programs and that the exercise was performed 2 or 3 times a week. Studies utilized a variety of modes; therefore, regardless of the equipment available, Mrs. Smith's physical therapist should be able to effect similar changes in her impairment at the body structure and function level, as well as her activity and participation levels, utilizing supervised PRT.

What can be advised based on the results of this systematic review?

Older adult patients, such as Mrs. Smith, are likely to benefit from supervised PRT regardless of the setting or mode of training. Based on the results of the Cochrane review, patients who participate in PRT are likely to show meaningful strength gains, along with decreases in physical disability. Substantial improvements in key functional activities including ADLs, gait speed, and the ability to get out of a chair should be expected. Larger changes in strength should be expected with higher intensities of PRT and in individuals without health problems or functional limitations.

References

1. [↵](#)
The Cochrane Library. Available at: <http://www.thecochranelibrary.com/view/0/index.html>. Accessed November 2, 2010.
2. [↵](#)
 1. Goodpaster BH,
 2. Park SW,
 3. Harris TB,
 4. et al

. *The loss of skeletal muscle strength, mass, and quality in older adults: the health, aging and body composition study. J Gerontol A Biol Sci Med Sci. 2006;61:1059-1064.*

[Abstract/FREE Full Text](#)
3. [↵](#)
 1. Rolland Y,
 2. Czerwinski S,
 3. Abellan Van Kan G,
 4. et al

. *Sarcopenia: its assessment, etiology, pathogenesis, consequences and future perspectives. J Nutr Health Aging. 2008;12:433-450.*

[CrossRefMedline](#)
4. [↵](#)
 1. Reid KF,
 2. Naumova EN,
 3. Carabello RJ,
 4. et al

. *Lower extremity muscle mass predicts functional performance in mobility-limited elders. J Nutr Health Aging. 2008;12:493-498.*

[CrossRefMedline](#)
5. [↵](#)
 1. Cruz-Jentoft AJ,
 2. Landi F,
 3. Topinková E,
 4. Michel JP

. *Understanding sarcopenia as a geriatric syndrome. Curr Opin Clin Nutr Metab Care. 2010;13:1-7.*

[CrossRefMedline](#)
6. [↵](#)
 1. Fried LP,
 2. Tangen CM,
 3. Walston J,
 4. et al.

; *Cardiovascular Health Study Collaborative Research Group. Frailty in older adults: evidence for a phenotype. J Gerontol A Biol Sci Med Sci. 2001;56:M146-M156.*

[Abstract/FREE Full Text](#)

7. [↵](#)

1. Newman AB,
2. Kupelian V,
3. Visser M,
4. et al.

. *Health ABC Study Investigators. Sarcopenia: alternative definitions and associations with lower extremity function. J Am Geriatr Soc. 2003;51:1602-1609.*

[CrossRefMedline](#)

8. [↵](#)

1. Janssen I,
2. Baumgartner RN,
3. Ross R,
4. et al

. *Skeletal muscle cutpoints associated with elevated physical disability risk in older men and women. Am J Epidemiol. 2004;159:413-421.*

[Abstract/FREE Full Text](#)

9. [↵](#)

1. Iannuzzi-Sucich M,
2. Prestwood KM,
3. Kenny AM

. *Prevalence of sarcopenia and predictors of skeletal muscle mass in healthy, older men and women. J Gerontol A Biol Sci Med Sci. 2002;57:M772-M777.*

[Abstract/FREE Full Text](#)

10. [↵](#)

1. Williams MA,
2. Haskell WL,
3. Ades PA,
4. et al.

. *American Heart Association Council on Clinical Cardiology; American Heart Association Council on Nutrition, Physical Activity, and Metabolism. Resistance exercise in individuals with and without cardiovascular disease: 2007 update: a scientific statement from the American Heart Association Council on Clinical Cardiology and Council on Nutrition, Physical Activity, and Metabolism. Circulation. 2007;116:572-584.*

[Abstract/FREE Full Text](#)

11. [↵](#)

1. Latham N,
2. Anderson C,
3. Bennett D,
4. Stretton C

. *Progressive resistance strength training for physical disability in older people. Cochrane Database Syst Rev. 2003;(2):CD002759.*

12. [↵](#)

1. Liu CJ,
2. Latham NK

. *Progressive resistance strength training for improving physical function in older adults. Cochrane Database Syst Rev. 2009;(3):CD002759.*

13.

1. Perera S,
2. Mody SH,
3. Woodman RC,
4. Studenski SA

. *Meaningful change and responsiveness in common physical performance measures in older adults. J Am Geriatr Soc. 2006;54:743-749.*

[CrossRefMedline](#)

14.

1. Kennedy DM,
2. Stratford PW,
3. Wessel J,
4. et al

. *Assessing stability and change of four performance measures: a longitudinal study evaluating outcome following total hip and knee arthroplasty. BMC Musculoskelet Disord. 2005;6:3.*

[CrossRefMedline](#)

15.

1. Mangione KK,
2. Craik RL,
3. McCormick AA,
4. et al

. *Detectable changes in physical performance measures in elderly African Americans. Phys Ther. 2010;90:921-927.*

[Abstract/FREE Full Text](#)

16. [↵](#)

1. Podsiadlo D,
2. Richardson S

. *The timed "Up & Go": a test of basic functional mobility for frail elderly persons. J Am Geriatr Soc. 1991;39(2):142-148.*

[Medline](#)

17. [↵](#)

1. Leiper CI,
2. Craik RL

. *Relationships between physical activity and temporal-distance characteristics of walking in elderly women. Phys Ther. 1991;71:791-803.*

[Abstract/FREE Full Text](#)

18. [↵](#)

Delmarva Foundation. Fall risk assessment form. Lincoln, Nebraska: CIMRO of Nebraska. Available at:

<http://www.bluefieldstate.edu/Schools/nursradt/Fall%202010/AS%20NURSING/NURS%20117L/Appendix%20B%20Pages%2039-41%20Fall%20Risk%20Assessment.pdf>.

Accessed November 8, 2010.